

Bounds of automorphism groups of genus 2 fibrations

Zhi-Jie Chen*

(Second version)
March 3, 1993

It is well known that the automorphism group of a surface of general type is finite and bounded by a function of K^2 [1]. Since then, several authors worked on this subject and found better upper bounds of the group. Recently Xiao [11,12] has obtained a linear bound for this group. Hence it is natural to investigate the upper bounds for particular classes of surfaces. Here we are interested in the upper bounds of various automorphism groups of surfaces with genus 2 pencils. As a first step, in the present paper, we will study the upper bounds of automorphism groups of genus 2 fibrations.

We always suppose that S is a smooth projective surface over the complex number field. A genus 2 fibration is a morphism $f : S \longrightarrow C$ where C is a projective curve such that a general fiber of f is a smooth curve of genus 2.

Definition 0.1 *An automorphism of the fibration $f : S \longrightarrow C$ is a pair of automorphisms $(\tilde{\sigma}, \sigma)$ where $\tilde{\sigma} \in \text{Aut}(S)$, $\sigma \in \text{Aut}(C)$, such that the diagram*

$$\begin{array}{ccc} S & \xrightarrow{\tilde{\sigma}} & S \\ f \downarrow & & \downarrow f \\ C & \xrightarrow{\sigma} & C \end{array}$$

*This work is realized under the support of NSF grant #DMS 9022140 while the author is visiting Mathematical Sciences Research Institute. Also partly supported by NSFC and K. C. Wong Education Foundation.

commutes.

The automorphism group of fibration f will be denoted by $\text{Aut}(f)$. The main results of this paper are the following

Theorem 0.1 *Suppose S is a surface of general type over the complex number field with a relatively minimal genus 2 fibration $f : S \longrightarrow C$. Then*

$$|\text{Aut}(f)| \leq 504K_S^2.$$

If f is not locally trivial, then

$$|\text{Aut}(f)| \leq 288K_S^2.$$

In particular,

$$|\text{Aut}(f)| \leq \begin{cases} 126K_S^2, & \text{if } g(C) \geq 2; \\ 144K_S^2, & \text{if } g(C) = 1; \\ 120K_S^2 + 960, & \text{if } g(C) = 0. \end{cases}$$

And these are all the best bounds.

Theorem 0.2 *Suppose S is a surface of general type over the complex number field with a relatively minimal genus 2 fibration $f : S \longrightarrow C$, G is an abelian automorphism group of f . Then*

$$|G| \leq 12.5K_S^2 + 100.$$

This bound is the best.

Theorem 0.3 *Suppose S is a surface of general type over the complex number field with a relatively minimal genus 2 fibration $f : S \longrightarrow C$, G is a cyclic automorphism group of f . Then*

$$|G| \leq \begin{cases} 5K_S^2, & \text{if } g(C) = 1, K_S^2 \geq 12; \\ 12.5K_S^2 + 90, & \text{if } g(C) = 0. \end{cases}$$

These bounds are the best.

Theorem 0.4 *Suppose S is a minimal surface of general type over the complex number field with a genus 2 fibration $f : S \longrightarrow C$ such that $g(C) \geq 2$, G is a cyclic automorphism group of f . Then*

$$|G| \leq 5K_S^2 + 30.$$

Theorem 0.1 will be obtained as a consequence of several propositions in Section 3. In Section 4, we discuss abelian and cyclic automorphism groups of the fibration f . The propositions proved there imply Theorems 0.2, 0.3 and 0.4. We remark that Xiao [7] has obtained a bound for abelian automorphism groups of f . Our theorem is an improvement of his. Examples are given in Section 5 to show that most of these bounds are the best possible.


1 Preliminaries

The surfaces with genus 2 pencils have been largely studied by many authors. The facts we needed in this paper mostly appeared in [3, 6, 9, 10]. In particular, Xiao's book [10] gave a systematic description of the properties of genus 2 fibrations which are just what we needed here. Unfortunately, this book has not been translated into English yet, hence it is not available for most readers. For this reason, we will recall some materials in this section.

Let $f : S \longrightarrow C$ be a relatively minimal fibration of genus 2, $\omega_{S/C} = \omega_S \otimes f^* \omega_C^\vee$ the relative canonical sheaf of f . For a sufficiently ample invertible sheaf $\mathcal{L}$ on C , the natural morphism $f^*(f_* \omega_{S/C} \otimes \mathcal{L}) \longrightarrow \omega_{S/C} \otimes f^* \mathcal{L}$ defines a natural map Φ :

$$\begin{array}{ccc} S & \xrightarrow{\quad \Phi \quad} & P = \mathbf{P}(f_* \omega_{S/C} \otimes \mathcal{L}) \\ f \searrow & & \swarrow \pi \\ & C & \end{array}$$

Φ is called a relatively canonical map. By a succession of blow-ups, we can obtain the following commutative diagram:



where ρ and ψ are compositions of finitely many blow-ups, $\tilde{\theta}$ is a double cover. Then we get the branch loci $\tilde{R}$ on $\tilde{P}$ and R on P such that $\tilde{R}$ is the minimal even resolution of R (i.e. the canonical resolution of double cover). If $\mathcal{L}$ is sufficiently ample, then all the singularities of R must be located in one of the 6 types of singular fibers defined by Horikawa [3]—0), I), II), III), IV) or V).

P is a relatively minimal ruled surface. We denote a section which has the least self-intersection by C_0 such that $C_0^2 = -e$. We will use F to denote both the fiber of f or of π .

A singular point of the branch locus will be called *negligible* if this point itself and all its infinitely near points are double points or triple points with at least 2 different tangents. By minimal even resolution, the inverse image of a negligible singular point is composed of (-2) -curves. All other singular points are called *non-negligible*. The singular fiber of type 0) in the classification of Horikawa is nothing else but the fiber which does not contain any non-negligible singular points.

The minimal even resolution $\psi : \tilde{P} \longrightarrow P$ can be decomposed into $\tilde{\psi} : \tilde{P} \longrightarrow \hat{P}$ followed by $\hat{\psi} : \hat{P} \longrightarrow P$, where $\tilde{\psi}$ and $\hat{\psi}$ are composed respectively of negligible and non-negligible blow-ups. The image of $\tilde{R}$ in $\hat{P}$ is denoted by $\hat{R}$.

If we take away all the isolated vertical (-2) -curves from the reduced divisor $\hat{R}$, we get a new reduced divisor $\hat{R}_p$, which is called the *principal part* of the branch locus $\hat{R}$. Then for any fiber F of $\pi : P \longrightarrow C$, the second and third *singularity index* of F , $s_2(F)$, $s_3(F)$, will be defined as follows:

If R has no quadruple singularities on F , then $s_3(F)$ equals the number of $(3 \rightarrow 3)$ type singularities of R on F . Otherwise $s_3(F)$ equals the number

of $(3 \rightarrow 3)$ type singularities of R on F plus 1. Hence $s_3(F) = 0$ if and only if R has no non-negligible singularities on F .

Let $\varphi : \hat{R}_p \rightarrow C$ be the natural projection induced by $\pi \circ \hat{\psi} : \hat{P} \rightarrow C$. Then the second singularity index $s_2(F)$ of F will be the ramification index of the divisor $\hat{R}_p$ on $f(F)$ with respect to the projection φ . If $\hat{R}_p$ has singularities (which must be negligible) on F , the singularity index $s_2(F)$ can be calculated as follows.

For a smooth point $p \in \hat{R}_p \cap F$, the ramification index of φ at p can be defined as for an ordinary smooth curve. If $p \in \hat{R}_p \cap F$ is a singular point of $\hat{R}_p$, then the ramification index of φ at p will be defined as the sum of ramification indices of the normalization of $\hat{R}_p$ at the preimage of p with respect to its projection to C plus the double of the influence to the arithmetic genus of $\hat{R}_p$ during its normalization at the singular point p . If the normalization of $\hat{R}_p$ contains an isolated vertical component E , then the contribution of E to the ramification index of φ is equal to $2g(E) - 2$.

As there are finite number of fibers F with $s_i(F) \neq 0$, we define the i -th *singularity index* of f , $s_i(f)$, to be the sum of $s_i(F)$ for all fibers, when $i = 2, 3$. If we take away from the branch locus R all the fibers F with odd $s_3(F)$, we obtain a divisor R_p which is called the *principal part* of R . Suppose that

$$R_p \sim -3K_{P/C} + nF,$$

where $K_{P/C}$ is the relative canonical divisor of π and $\sim$ represents numerical equivalence. With these definitions, the formula computing the relative invariants of a genus 2 fibration can be stated as follows.

Theorem 1.1 (Xiao [10]) *Let $f : S \rightarrow C$ be a relatively minimal fibration of genus 2. Then*

$$\begin{aligned} K_{S/C}^2 &= K_S^2 - 8(g(C) - 1) = \frac{1}{5}s_2(f) + \frac{7}{5}s_3(f) = 2n - s_3(f), \\ \chi_f &= \chi(\mathcal{O}_S) - (g(C) - 1) = \frac{1}{10}s_2(f) + \frac{1}{5}s_3(f) = n - s_3(f). \end{aligned}$$

2 Local cases

We begin with a local fibration $f : S_\Delta \rightarrow \Delta$ where f is an analytic mapping onto the unit disk Δ , S_Δ is a 2-dimensional analytic smooth manifold and

the fibers of f are projective curves. We assume that the fiber of the zero is singular and all the fibers over $\Delta^* = \Delta - \{0\}$ are smooth curves of genus 2.

Similarly, we have a commutative diagram

$$\begin{array}{ccc}
 \tilde{S}_\Delta & \xrightarrow{\tilde{\theta}} & \tilde{P}_\Delta \\
 \rho \downarrow & & \downarrow \psi \\
 S_\Delta & \xrightarrow{\Phi} & P_\Delta \\
 f \searrow & & \swarrow \pi \\
 & \Delta &
 \end{array}$$

Denote the branch locus in P_Δ by R_Δ . We also denote the horizontal part of R_Δ by R'_Δ , that is,

$$R'_\Delta = \begin{cases} R_\Delta - F_0, & \text{if } R_\Delta \text{ contains } F_0, \\ R_\Delta & \text{otherwise.} \end{cases}$$

Let $F_0 = \pi^{-1}(0)$, $F_t = \pi^{-1}(t)$, $t \in \Delta^*$, and $K_\Delta = \{\tilde{\sigma} \in \text{Aut}(S_\Delta) | f \circ \tilde{\sigma} = f\}$. Any automorphism $\tilde{\sigma} \in K_\Delta$ induces an automorphism σ of P_Δ satisfying $\pi \circ \sigma = \pi$ and $\sigma(R_\Delta) = R_\Delta$. We denote the image of K_Δ by $\bar{K}_\Delta \subseteq \text{Aut } P_\Delta$, then

$$|K_\Delta| = 2|\bar{K}_\Delta|.$$

Note that any finite automorphism subgroup of $\mathbf{P}^1$ must be one of the following:

$G \subseteq \text{Aut}(\mathbf{P}^1)$		$ G $	Number of points in an orbit
Cyclic group	Z_n	n	1, n
Dihedral group	D_{2n}	$2n$	2, n , $2n$
Tetrahedral group	T_{12}	12	4, 6, 12
Octahedral group	O_{24}	24	6, 8, 12, 24
Icosahedral group	I_{60}	60	12, 20, 30, 60

For any $\sigma \in \bar{K}_\Delta$, its restriction to $F_t \cong \mathbf{P}^1$, $\sigma|_{F_t}$, must preserve the set of 6 points contained in $F_t \cap R_\Delta$. Hence $\bar{K}_\Delta$ can be isomorphic to the following groups: O_{24} , T_{12} , D_{12} , D_6 , Z_6 , Z_5 , D_4 , Z_4 , Z_3 , Z_2 and 1.

Lemma 2.1 *If $\bar{K}_\Delta \cong O_{24}, T_{12}$ or D_{12} , then F_0 is contained in R_Δ and R_Δ has 6 ordinary double points on F_0 . In this case, we have $s_2(F_0) = 10$, $s_3(F_0) = 0$.*

Proof. Since $\bar{K}_\Delta \cong O_{24}, T_{12}$ or D_{12} , $R_\Delta \cap F_t$ ($t \in \Delta^*$) consists respectively of 6 vertices of a regular octahedron, of 6 points corresponding to the centers of edges of a regular tetrahedron, or of sixth roots of unit. These 6 horizontal branches of R_Δ cannot intersect when $t \rightarrow 0$. But by assumption, R_Δ must have some singularities, so F_0 is contained in R_Δ .

Since R_Δ does not contain non-negligible singularities, one has $s_3(F_0) = 0$ and $R_\Delta = \hat{R}_\Delta = (\hat{R}_\Delta)_p$. On F_0 , R_Δ has 6 ordinary double points, the influence of each double point to the arithmetic genus of R_Δ during its normalization is equal to 1. The preimage of F_0 in the normalization of R_Δ is a smooth vertical rational curve which does not meet any other branches, so its contribution to the index $s_2(F_0)$ is equal to -2 . Therefore $s_2(F_0) = 2 \times 6 + (-2) = 10$. $\square$

We list the following useful lemmas, the proof is evident. Since local equations are used for calculation of singularity indices, they are given in simplified form, omitting some higher order terms. All the non-negligible singularities here are canonical, i.e. defined by Horikawa.

Lemma 2.2 *If $\bar{K}_\Delta \cong D_6$ and R'_Δ is not étale over Δ , then up to a coordinates transformation,*

(1) *The equation of R'_Δ is $(x^3 - t^k)(t^k x^3 - 1)$, $k > 0$. In this case, $s_3(F_0) = 0$ implies $s_2(F_0) \geq 4$.*

(2) *The equation of R'_Δ is $(x^3 - 1)^2 - t^k(x^3 + 1)^2$, $k > 0$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 3$.*

Lemma 2.3 *If $\bar{K}_\Delta \cong Z_6$ and R'_Δ is not étale over Δ , then up to a coordinates transformation, the equation of R'_Δ is $x^6 - t^k$, $1 \leq k \leq 3$. If $k = 3$, it has a non-negligible singularity and $s_3(F_0) = 1$, $s_2(F_0) = 3$. Otherwise $s_2(F_0) \geq 5$.*

Lemma 2.4 *If $\bar{K}_\Delta \cong Z_5$ and R'_Δ is not étale over Δ , then up to a coordinates transformation,*

(1) The equation of R'_Δ is $x(x^5 - t^k)$, $k = 1, 2$. In this case, $s_3(F_0) = 0$, $s_2(F_0) \geq 6$.

(2) The equation of R'_Δ is $x(t^k x^5 - 1)$, $k = 1, 2$. In this case, $s_3(F_0) = 0$, $s_2(F_0) \geq 4$.

Lemma 2.5 *If $\bar{K}_\Delta \cong D_4$ and R'_Δ is not étale over Δ , then up to a coordinates transformation,*

(1) *The equation of R'_Δ is $(x^2 - 1)((x - 1)^2 - t^k(x + 1)^2)(t^k(x - 1)^2 - (x + 1)^2)$, $k > 0$. In this case, $s_3(F_0) = 0$ implies $s_2(F_0) \geq 6$.*

(2) *The equation of R'_Δ is $(x^2 - 1)(x^2 - t^k)(t^k x^2 - 1)$, $k > 0$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 2$.*

Lemma 2.6 *If $\bar{K}_\Delta \cong Z_4$ and R'_Δ is not étale over Δ , then up to a coordinates transformation, the equation of R'_Δ is $x(x^4 - t^k)$, $k = 1, 2$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 5$.*

Lemma 2.7 *If $\bar{K}_\Delta \cong Z_3$ and R'_Δ is not étale over Δ , then up to a coordinates transformation,*

(1) *The equation of R'_Δ is $(x^3 - t^{k_1})(t^{k_2} x^3 - a(t))$, $k_1, k_2 > 0$, $a(0) \neq 0$. In this case, $s_3(F_0) = 0$ implies $s_2(F_0) \geq 4$.*

(2) *The equation of R'_Δ is $x^6 + a(t)x^3 + t^k$, $1 \leq k \leq 3$. In this case, $s_3(F_0) = 0$ implies $s_2(F_0) \geq 5$.*

(3) *The equation of R'_Δ is $(x^3 - b - t^{k_1})(x^3 - b - t^{k_2} a(t))$, $k_1, k_2 > 0$, $a(0) \neq 0$ and $b \neq 0$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 6$.*

(4) *The equation of R'_Δ is $(x^3 - t^k)(x^3 - a(t))$, $1 \leq k \leq 3$, $a(0) \neq 0$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 2$.*

(5) *The equation of R'_Δ is $((x - b)^2 - t^k a(t))((x - b\omega)^2 - \omega^2 t^k a(t))((x - b\omega^2)^2 - \omega t^k a(t))$, $k > 0$, $a(0) \neq 0$, $b \neq 0$, $\omega = \exp(2\pi i/3)$. In this case, we have $s_3(F_0) = 0$, $s_2(F_0) \geq 3$.*

We summarize the results of Lemmas 2.2 through 2.7 in the following table. Here we assume that R'_Δ has only negligible singularities or ramifications on F_0 .

$\bar{K}_\Delta$	$ K_\Delta $	$s_2(F_0)$	$ K_\Delta /s_2(F_0)$
D_6	12	≥ 3	≤ 4
Z_6	12	≥ 5	≤ 2.4
Z_5	10	≥ 4	≤ 2.5
D_4	8	≥ 2	≤ 4
Z_4	8	≥ 5	≤ 1.6
Z_3	6	≥ 2	≤ 3
Z_2	4	≥ 1	≤ 4
1	2	≥ 1	≤ 2

Lemma 2.8 *If R'_Δ has only negligible singularities or ramifications on F_0 , then $|K_\Delta|/s_2(F_0) \leq 4$. Moreover, if $\bar{K}_\Delta \cong Z_6, Z_5, Z_4$ or 1, then $|K_\Delta|/s_2(F_0) \leq 2.5$.*

3 Bounds of automorphism groups

Let $G = \text{Aut}(f)$ be the automorphism group of the fibration of genus two $f : S \longrightarrow C$. Then we have an exact sequence

$$1 \longrightarrow K \longrightarrow G \longrightarrow H \longrightarrow 1,$$

$$(\tilde{\sigma}, \sigma) \mapsto \sigma$$

where $H \subseteq \text{Aut}(C)$, $K = \{(\tilde{\sigma}, \text{id}) \in G\} = \{\tilde{\sigma} \in \text{Aut}(S) | f \circ \tilde{\sigma} = f\}$. Thus

$$|G| = |K||H|.$$

The elements of H are often regarded as transformations of the fibers of f or π .

Proposition 3.1 *If $f : S \longrightarrow C$ is a relatively minimal fibration of genus 2 with $g(C) \geq 2$, then*

$$|\text{Aut}(f)| \leq 504K_S^2.$$

Proof. Since $|K| \leq 48$, $|H| \leq |\text{Aut}(C)| \leq 84(g(C) - 1)$, we have

$$|G| = |K||H| \leq 4032(g(C) - 1).$$

On the other hand, $K_{S/C}^2 \geq 0$ and the equality holds if and only if f is locally trivial. Hence

$$K_S^2 \geq 8(g-1)(g(C)-1) = 8(g(C)-1),$$

and

$$|G| \leq 504K_S^2. \quad \square$$

Proposition 3.2 *If $f : S \longrightarrow C$ is a relatively minimal fibration of genus 2 which is not locally trivial with $g(C) \geq 2$, then*

$$|\text{Aut}(f)| \leq 126K_S^2.$$

Proof. Let R' denote the horizontal part of the branch locus R . If R' is not étale over C , then by Lemmas of Section 2, we have $|K| \leq 12$. As $|H| \leq 84(g(C)-1) \leq 10.5K_S^2$,

$$|G| \leq 12|H| \leq 126K_S^2.$$

Now assume that R' is étale. Since f is not locally trivial, we must have $K_{S/C}^2 > 0$, i.e. either $s_3(f) > 0$ or $s_2(f) > 0$. So R must contain some fiber F_0 . By Lemma 2.1, $s_3(F_0) = 0$, $s_2(F_0) = 10$. Let $p = f(F_0)$, $n = |H|$. Since H is a subgroup of $\text{Aut}(C)$, H determines a finite morphism $\tau : C \longrightarrow X = C/H$. Denote the ramification index of $p \in C$ with respect to τ by r and the other ramification indices by r_i . Then the Hurwitz's theorem implies that

$$2g(C) - 2 = n(2g(X) - 2) + n \sum \left(1 - \frac{1}{r_i}\right).$$

As the H -orbit of the point p has n/r points, this implies that $s_2(f) \geq 10n/r$. Hence

$$\begin{aligned} K_S^2 &\geq \frac{1}{5}s_2(f) + 8(g(C)-1) = \frac{2n}{r} + 4n \left[2g(X) - 2 + \sum \left(1 - \frac{1}{r_i}\right)\right] \\ &= 4n \left[2g(X) - 2 + \frac{1}{2r} + \sum \left(1 - \frac{1}{r_i}\right)\right]. \end{aligned}$$

It is not difficult to see that the expression $2g(X) - 2 + 1/2r + \sum(1 - 1/r_i)$ reaches its minimal value $2/21$ (under the condition $2g(X) - 2 + \sum(1 - 1/r_i) > 0$) when $g(X) = 0$, $r_1 = 2$, $r_2 = 3$, and $r = r_3 = 7$. That is

$$K_S^2 \geq \frac{8}{21}n = \frac{8}{21}|H|.$$

Thus

$$|G| \leq 48|H| \leq 126K_S^2. \quad \square$$

Remark. It is not difficult to see that if f is not locally trivial with $g(C) \geq 2$ and $|\text{Aut}(f)| = 126K_S^2$, then $|\text{Aut}(C)| = 84(g(C) - 1)$, $|\text{Aut}(F)| = 48$ for any smooth fiber F and $\text{Aut}(f) \cong \text{Aut}(C) \times \text{Aut}(F)$. We will give an example later. In this case, the fibration f is of constant moduli and $S/\text{Aut}(f) \cong F_1$.

Lemma 3.1 *Let S be a surface of general type which has a relatively minimal genus 2 fibration $f : S \rightarrow C$. If the third singularity index $s_3(f) \neq 0$, then*

$$|\text{Aut}(f)| \leq \frac{60}{7}rK_{S/C}^2,$$

where

$$r = \min_{s_3(F) \neq 0} |\text{Stab}_H f(F)|,$$

$\text{Stab}_H f(F)$ is the stabilizer of $f(F)$ in H .

Proof. Let F_0 be a singular fiber such that $s_3(F_0) \neq 0$ and $r = |\text{Stab}_H f(F_0)|$. Then

$$K_{S/C}^2 \geq \frac{7}{5}s_3(f) \geq \frac{7s_3(F_0)}{5r}|H|,$$

and we get

$$|G| = |K||H| \leq \frac{r}{s_3(F_0)} \cdot \frac{60}{7}K_{S/C}^2 \leq \frac{60}{7}rK_{K/C}^2. \quad \square$$

Lemma 3.2 *Let S be a surface of general type which has a relatively minimal genus 2 fibration $f : S \rightarrow C$. If the horizontal part R' of the branch locus R is not étale and has only negligible singularities or ramifications, then*

$$|\text{Aut}(f)| \leq 20rK_{S/C}^2,$$

where

$$r = \min_{F \text{ singular fiber}} |\text{Stab}_H f(F)|.$$

Proof. Let F_0 be a singular fiber with $r = |\text{Stab}_H f(F_0)|$. Since here

$$K_{S/C}^2 \geq \frac{1}{5}s_2(f) \geq \frac{s_2(F_0)}{5r}|H|,$$

we have

$$|G| = |K||H| \leq \frac{r|K|}{s_2(F_0)} \cdot 5K_{S/C}^2 \leq 20rK_{S/C}^2,$$

by Lemma 2.8. $\square$

Lemma 3.3 *Let S be a surface of general type which has a relatively minimal genus 2 fibration $f : S \rightarrow C$. If the horizontal part R' of the branch locus R is étale, then*

$$|\text{Aut}(f)| \leq 24rK_{S/C}^2,$$

where

$$r = \min_{F \text{ singular fiber}} |\text{Stab}_H f(F)|.$$

Proof. Let F_0 be a singular fiber with $r = |\text{Stab}_H f(F_0)|$. By assumption, we have $s_2(F_0) = 10$. Hence

$$|G| = |K||H| \leq \frac{r|K|}{s_2(F_0)} \cdot 5K_{S/C}^2 \leq 24rK_{S/C}^2. \quad \square$$

Let $\bar{K}$ denote the subgroup in $\text{Aut}(P)$ which is induced by K . If $\sigma \in \bar{K}$, then $\pi \circ \sigma = \pi$ and $\sigma(R) = R$. Let K_1 be a cyclic subgroup of order m of $\bar{K}$, $Q = P/K_1$ be the quotient surface. Then Q is a ruled surface. We have a commutative diagram

$$\begin{array}{ccc} P & \xrightarrow{\alpha} & Q \\ \pi \searrow & & \swarrow \pi' \\ & C & \end{array}$$

Let C_0 and $C_\infty \sim C_0 + eF$ be the reduced ramification divisors of K_1 . Let C'_0 be a section of π' with the least self-intersection $C_0'^2 = -e'$, F' be a general fiber of π' . Then $\alpha^*C'_0 = mC_0$, $\alpha^*C'_\infty = mC_\infty$, $\alpha^*F' = F$ and $e' = me$. Let $D = \alpha(R')$, $C' = C'_0 + C'_\infty$ be the branch locus. Then $C' \sim 2C'_0 + e'F' \sim -K_{Q/C}$.

Lemma 3.4 *Assume $\bar{K} \cong D_6$. If R' is not étale and has only negligible singularities or ramifications, then f has more than one H -orbit of singular fibers.*

Proof. Let K_1 be the unique cyclic subgroup of order 3 of $\bar{K}$. There are 2 types of singular fibers as listed in Lemma 2.2. Let F_0 be a singular fiber. Then the local equations of D near F_0 are (1) $(x - t^k)(t^k x - 1)$, $k \leq 3$, (2) $(x - 1)^2 - t^k(x + 1)^2$, $k > 0$. In case (1), D meets C' at 2 points in F_0 . In case (2), D does not meet C' in F_0 .

If all the singular fibers of f is of type (1), then D is an étale cover of C . This means that $a = e'$, $C' \sim D$. Hence $DC' = 0$ which is impossible because D and C' meet in F_0 .

If all the singular fibers of f is of type (2), then $DC' = 0$. Hence $D \sim C'$ and $D(D + K_{Q/C}) = 0$. This means that D is étale over C . A contradiction. $\square$

Lemma 3.5 *Assume $\bar{K} \cong D_4$. If R' is not étale, then f has more than one H -orbit of singular fibers.*

If H is cyclic and $g(C) = 0$, then

$$|\text{Aut}(f)| \leq 12.5K_{S/C}^2.$$

Proof. In this case, there are 4 sections in P which do not meet each other. Hence $e = 0$. R' contains 2 of these sections and denoted by C_0 and C_∞ . Let K_1 be a cyclic subgroup of $\bar{K}$ with C_0 and C_∞ as ramifications. Assume that there is only one H -orbit of singular fibers. If these singular fibers are all of type (1) in Lemma 2.5, then the local equation of $D = \alpha(R' - C_0 - C_\infty)$ is $(x - t^k)(t^k x - 1)$, namely, D is étale. Therefore $D \sim 2C'_0$, $DC'_0 = DC'_\infty = 0$, a contradiction. If the singular fibers are of type (2) in Lemma 2.5, then D does not meet C'_0 and C'_∞ . Hence $D \sim 2C'_0$, $D^2 = 0$, a contradiction. That implies there are at least 2 H -orbits.

Now suppose H is cyclic. Let $h = |H|$. We call an H -orbit *big* if it contains h fibers. If there is a big H -orbit whose singular fibers are of type (1), then $s_2(F_0) \geq 6$, so $|G| \leq (20/3)K_{S/C}^2$. If $|G| > (20/3)K_{S/C}^2$, then the singular fibers in a big H -orbit must be of type (2) with $k \leq 2$. Let F_2 and F_3 denote 2 fibers fixed by H . Then at least one of them is of type (1). The structure of types (1) and (2) implies that the normalization of $D = \alpha(R' - C_0 - C_\infty)$ is étale with respect to π' . Hence D must decompose into 2 isomorphic sections D_1 and D_2 , $D_1 \sim D_2 \sim C'_0 + aF'$. Since both

D_1 and D_2 meet C'_0 and C'_∞ , F_2 and F_3 are all singular of type (1). Since $D_1 D_2 = 2a = kh$, $D_1 C'_0 = a = kh/2$. Hence the local equation of R' near F_2 or F_3 is $(x^2 - 1)((x - 1)^2 - t^{kh/2}(x + 1)^2)(t^{kh/2}(x - 1)^2 - (x + 1)^2)$. When $h \geq 6$, these are non-negligible singularities. If F_i ($i = 2, 3$) is singular fiber of type I), $s_3(F_i) = 2[(kh - 2)/8] + 1 \geq (kh - 1)/4$. If F_i is of type II), $s_3(F_i) = 2[kh/8] \geq (kh - 6)/4$. So

$$K_{S/C}^2 \geq \frac{1}{5} \times 2 \times h + \frac{7}{5} \times \frac{h - 6}{4} \times 2 = \frac{11}{10}h - \frac{21}{5}.$$

$$|G| = 8h \leq \frac{80}{11}(K_{S/C}^2 + \frac{21}{5}) < 12.5K_{S/C}^2.$$

If there are more than one big H -orbits, it can be similarly shown that $|G| \leq 12.5K_{S/C}^2$. $\square$

Lemma 3.6 *Assume $\bar{K} \cong Z_3$. If R' is not étale and has only negligible singularities or ramifications and f has only one H -orbit of singular fibers, then*

$$|\text{Aut}(f)| \leq 6rK_{S/C}^2.$$

where

$$r = \min_{F \text{ singular fiber}} |\text{Stab}_H f(F)|.$$

Proof. Let $K_1 = \bar{K}$. If the singular fibers are of types (1) or (4) in Lemma 2.7, then $D \sim 2C'_0 + aF'$ is étale. $D(K_{Q/C} + D) = 0$ implies $a = e'$. Hence $D(C'_0 + C'_\infty) = 0$, a contradiction. If the singular fiber F_0 is of type (5) with $k = 1$, then D is irreducible and smooth near F_0 . This implies $DC'_\infty \neq 0$, a contradiction. Therefore $s_2(F_0) \geq 5$ for any singular fiber F_0 . So $|G| \leq 6rK_{S/C}^2$. $\square$

Lemma 3.7 *Assume $\bar{K} \cong Z_2$. If R' is not étale and f has only one H -orbit of singular fibers, then*

$$|\text{Aut}(f)| \leq 5rK_{S/C}^2.$$

where

$$r = \min_{F \text{ singular fiber}} |\text{Stab}_H f(F)|.$$

Proof. Let F_0 be a singular fiber. $|G| > 5rK_{S/C}^2$ implies $s_2(F_0) \leq 3$. We distinguish between 2 cases.

Case I. R' contains C_0 and C_∞ . Then the local equation of R' near F_0 must be (1) $x(x^2 - t)(x^2 - a(t))$, $a(0) \neq 0$, $s_2(F_0) = 3$; (2) $x((x^2 - a^2)^2 - t)$, $a \neq 0$, $s_2(F_0) = 2$. Let $D = \alpha(R' - C_0 - C_\infty) \sim 2C'_0 + aF'$. If all the singular fibers are of type (1), then D is étale. This is impossible. If the singular fibers are of type (2), then D is irreducible and does not meet C' . This is impossible.

Case II. R' does not contain C_0 and C_∞ . Then the local equation of R' may be (1) $(x^2 - t)(x^2 - a(t))(x^2 - b(t))$, $a(0)b(0) \neq 0$, $a(0) \neq b(0)$, $s_2(F_0) = 1$; (2) $(x^2 - t)(ta(t)x^2 - 1)(x^2 - b(t))$, $a(0)b(0) \neq 0$, $s_2(F_0) = 2$; (3) $((x^2 - a^2)^2 - t)(x^2 - b(t))$, $ab(0) \neq 0$, $s_2(F_0) = 2$; (4) $((x^2 - a^2)^2 - t)(x^2 - tb(t))$, $b(0) \neq 0$, $s_2(F_0) = 3$. Let $D = \alpha(R') \sim 3C'_0 + aF'$. If F_0 is of type (1) or (2), then D is étale and smooth. D must be decomposed into 3 disjoint components. This means $e' = 0$, a contradiction. If F_0 is of type (3) or (4), then D is smooth. The ramification index $D(D + K_{Q/C}) = 4a - 6e' = |H|/r$. Hence $DC' = 2a - 3e' = |H|/2r$. This is a contradiction because we have $DC' = 0$ for type (3) and $DC' = |H|/r$ for type (4). $\square$

Proposition 3.3 *If S is a minimal surface of general type which has a genus 2 fibration $f : S \rightarrow C$ with $g(C) = 1$, then*

$$|\text{Aut}(f)| \leq 144K_S^2.$$

Proof. In this case, we have

$$K_S^2 = K_{S/C}^2 = \frac{1}{5}s_2(f) + \frac{7}{5}s_3(f) > 0.$$

Thus either $s_3(f) > 0$ or $s_2(f) > 0$.

Let $j(C)$ be the j -invariant of the elliptic curve C . Let m denote the number of points contained in a smallest H -orbit of C . Since H is a finite subgroup of $\text{Aut}(C)$, we have

$$m = \begin{cases} |H|/2 & \text{if } j(C) \neq 0, 1728, \\ |H|/4 & \text{if } j(C) = 1728, \\ |H|/6 & \text{if } j(C) = 0. \end{cases}$$

Since $r \leq 6$, by Lemma 3.1, 3.2 and 3.3, the conclusion is immediate. $\square$

Proposition 3.4 *If S is a surface of general type which has a relatively minimal fibration of genus two $f : S \longrightarrow C$ with $g(C) = 0$, then*

$$|\mathrm{Aut}(f)| \leq 120(K_S^2 + 8).$$

Moreover, we have

$$|\mathrm{Aut}(f)| \leq 48(K_S^2 + 8)$$

for $K_S^2 \geq 33$, and when $K_S^2 \leq 32$, there are only 4 exceptions.

Proof. In this case, we have

$$K_S^2 + 8 = K_{S/C}^2 = \frac{1}{5}s_2(f) + \frac{7}{5}s_3(f) > 0.$$

Hence either $s_3(f) > 0$ or $s_2(f) > 0$.

Case I. Assume that R' is étale over C . If $r \leq 5$, then by Lemma 3.3

$$|G| \leq 24rK_{S/C}^2 \leq 120(K_S^2 + 8).$$

If $r \geq 6$, then H must be cyclic or dihedral group. In this case, there are at most 2 singular fibers. Hence $K_{S/C}^2 \leq 4$ by Theorem 1.1. This means S is not of general type [10, Theorem 4.2.5, p.90].

Case II. Assume that R' is not étale. Then f is a fibration of variable moduli. Hence f must contain more than 2 singular fibers ([2]). This implies $r \leq 5$. The conclusion is obtained by Lemmas 3.1 and 3.2.

In the preceding argument, we can see that $|G| \leq 48(K_S^2 + 8)$ holds if $r \leq 2$. If $|G| > 48(K_S^2 + 8)$, we must have $r > 3$. Then H is one of T_{12} , O_{24} or I_{60} .

If f has more than one H -orbit of singular fibers, then

$$\begin{aligned} \frac{K_{S/C}^2}{|G|} &\geq \frac{1}{5r} \left(\frac{s_2(F_0)}{|K|} + \frac{7s_3(F_0)}{|K|} \right) + \frac{1}{5r_1} \left(\frac{s_2(F_1)}{|K|} + \frac{7s_3(F_1)}{|K|} \right) \\ &\geq \frac{1}{25} \times \frac{1}{4} + \frac{1}{20} \times \frac{1}{4} = \frac{9}{400} > \frac{1}{48}. \end{aligned}$$

Therefore f has only one H -orbit.

If the singular fibers has non-negligible singularities, then by Lemma 3.1, $|G| \leq (60/7)rK_{S/C}^2 \leq (300/7)K_{S/C}^2 < 48K_{S/C}^2$. Suppose that the horizontal

part R' of the branch locus has only negligible singularities or ramifications, then by Lemmas 3.4, 3.5, 3.6 and 3.7, we have

$$|G| \leq 12.5rK_{S/C}^2.$$

Thus $|G| > 48K_{S/C}^2$ implies that $r \geq 4$ and $\bar{K}$ is Z_6 or Z_5 . If $\bar{K} \cong Z_6$, then $r = 5$ and $H \cong I_{60}$. To ensure $|G| > 48K_{S/C}^2$, we have $s_2(F_0) = 5$, i.e. $R = R' \sim -3K_{P/C} + nF$ is a smooth irreducible divisor. As a multiple cover on C , the ramification index of R is equal to $R(R + K_{P/C}) = 12n$. On the other hand, this ramification index is equal to $5 \times (60/5) = 60$, i.e. $n = 5$. But $2n = 10 = K_{S/C}^2 \neq s_2(f)/5 = 12$, a contradiction.

If $\bar{K} \cong Z_5$, then $|G| > 48K_{S/C}^2$ implies $s_2(F_0) = 4$. In this case $R = R' = C_0 + R_1$ where $R_1 \sim 5C_0 + (n + 3e)F$ is an smooth irreducible divisor and $R_1C_0 = 0$, i.e. $n = 2e$. Computing the ramification index of R_1 we get $R_1(R_1 + K_{P/C}) = 10n = 4|H|/r$. This implies $5r||H|$, a contradiction. Hence $|G| > 48(K_S^2 + 8)$ implies that R' is étale over C . There are only finite number of possibilities. We list the possible fibrations with $|G| > 48(K_S^2 + 8)$ as follows.

H	r	$ G $	K_S^2	$ K /(K_S^2 + 8)$	$ K /K_S^2$
I_{60}	5	2880	16	120	180
I_{60}	3	2880	32	72	90
O_{24}	4	1152	4	96	288
O_{24}	3	1152	8	72	144

In the Section 5 we will show their existence. $\square$

Corollary 3.5 *If S is a minimal surface of general type which has a genus 2 fibration $f : S \rightarrow C$ with $g(C) = 0$, then*

$$|\text{Aut}(f)| \leq 288K_S^2.$$

Proof. If $K_S^2 \geq 2$, then

$$48(K_S^2 + 8) < 288K_S^2.$$

By Proposition 3.4 we need only check the 4 exceptional examples. That fact leads to the inequality $|G| \leq 288K_S^2$. $\square$

4 Abelian automorphism groups

Let $G \subseteq \text{Aut}(f)$ be an abelian group. Then it is well known that $|K| \leq 12$.

Proposition 4.1 ([7, Lemma 8]) *Let $f : S \rightarrow C$ be a relatively minimal fibration of genus 2 with $g(C) \geq 2$, G is an abelian automorphism group of S , then*

$$|G| \leq 6K_S^2 + 96.$$

Let $\bar{G} \subseteq \text{Aut}(P)$ be the induced automorphism group of a commutative group G , then

$$1 \longrightarrow \bar{K} \longrightarrow \bar{G} \longrightarrow H \longrightarrow 1.$$

Lemma 4.1 *Assume that $\bar{K} \cong Z_3$, $g(C) = 0$. Let $p \in C$ be a fixed point of the cyclic group H , $F = \pi^{-1}(p)$. If there is a $\bar{K}|_F$ -orbit containing 3 points in F , then*

$$s_2(F) \geq 3|H|.$$

Proof. Since p is a fixed point of H , the induced action of $\bar{G}$ on F forms a commutative subgroup $\bar{G}|_F \subseteq \text{Aut}(F) \cong \text{Aut}(\mathbf{P}^1)$. Since $\bar{G}|_F$ stabilize this $\bar{K}|_F$ -orbit, $\bar{G}|_F = \bar{K}|_F \cong Z_3$, i.e. $H|_F = 1$. Hence the local equation of R' near F has the form $f(x^3, t^h)$ where $h = |H|$. Or explicetly, the local equation of R' are (3) $(x^3 - b - t^{k_1 h} a_1(t^h))(x^3 - b - t^{k_2 h} a_2(t^h))$; (5) $((x - b)^2 - t^{kh} a(t^h))((x - b\omega)^2 - \omega^2 t^{kh} a(t^h))((x - b\omega^2)^2 - \omega t^{kh} a(t^h))$, $b \neq 0$. Thus $s_2(F) \geq 3h = 3|H|$. $\square$

Proposition 4.2 *If S is a surface of general type which has a relatively minimal fibration of genus two $f : S \rightarrow C$ with $g(C) \leq 1$, G is an abelian automorphism group of f , then*

$$|G| \leq 12.5(K_S^2 + 8).$$

Proof. It is well known that H must be a cyclic group or a dihedral group $D_4 \cong Z_2 \oplus Z_2$.

If $g(C) = 1$ and that H does not act freely on C , then $|H| \leq 6$. Hence $|G| \leq 72 < 12.5(K_S^2 + 8)$. If $g(C) = 0$ and $H \cong D_4$, then $|G| \leq 48$, the claim holds too. So we can assume that H is a cyclic group and that there exists a singular fiber F_0 with $|\text{Stab}_H f(F_0)| = 1$.

Case I. Suppose that the horizontal part R' of the branch locus R is étale over C . Then

$$|G| \leq 6K_{S/C}^2.$$

Case II. Suppose that R' is not étale. If there is a big H -orbit with $s_3(F_0) \neq 0$, then

$$K_{S/C}^2 \geq \frac{7}{5}s_3(f) \geq \frac{7}{5}|H|,$$

so

$$|G| \leq \frac{60}{7}K_{S/C}^2 < 12.5(K_S^2 + 8).$$

Now suppose that on the big H -orbits R' has only negligible singularities or ramifications. If $\bar{K} \cong Z_6, Z_5, Z_4$ or 1, then by Lemma 2.8, we have

$$|G| \leq \frac{|K|}{s_2(F_0)} \cdot 5K_{S/C}^2 \leq 12.5K_{S/C}^2 \leq 12.5(K_S^2 + 8).$$

Suppose that $\bar{K} \cong D_4, Z_3$ or Z_2 and that $|G| > 12.5(K_S^2 + 8)$. Then Lemmas 3.5, 3.6 and 3.7 implies that f must have more than one H -orbit of singular fibers. To ensure $|G| > 12.5(K_S^2 + 8)$, f cannot have more than one big H -orbits. Thus we have $g(C) = 0$. Lemma 3.5 excludes the case of $\bar{K} \cong D_4$.

If $\bar{K} \cong Z_3$, then $s_2(F_0) \leq 2$. Hence F_0 must be of type (4) of Lemma 2.7 with $k = 1$. Taking $K_1 = \bar{K}$ we construct the quotient surface $Q = P/K_1$ as in §3. Then $D = \alpha(R')$ is étale near F_0 . But D cannot be étale. Hence at least one of the H -stabilized fibers F_2 or F_3 is of type (2) $k = 1$ or type (5) $k = 1$. Lemma 4.1 excludes the case of type (5). Suppose one of the F_i is of type (2). Then $D \sim 2C'_0 + aF'$ is irreducible and smooth. As a smooth double cover of $C \cong \mathbf{P}^1$, the ramification index of D is at least 2. So F_2 and F_3 are all of type (2). Then $DC' = D(D + K_{Q/C}) = 2(a - e') = 2$, a contradiction.

If $\bar{K} \cong Z_2$, then $s_2(F_0) = 1$. Hence the local equation of R' near F_0 is $(x^2 - t)(x^2 - a(t))(x^2 - b(t))$, $a(0)b(0) \neq 0$, $a(0) \neq b(0)$. So $D = \alpha(R')$ is étale near F_0 .

If F_2 and F_3 have no ramifications, the D can be decomposed into 3 components $D_i \sim C'_0 + a_iF'$, $i = 1, 2, 3$. These 3 components must meet each other on F_2 and F_3 . So there exists at least one point on F_i where 3 components intersect. The local equation of R' will be $(x^4 + a(t)x^2 + t^2)(x^2 - t^2b(t))$. But as $D_i(C'_\infty - C'_0) = e'$, we will have $|H| \leq 1$.

If F_2 or F_3 has ramifications, the equation of R' near F_i must be (1) $x^6 - t$; (2) $(x^4 - t)(t^k a(t)x^2 - 1)$, $a(0) \neq 0$; (3) $((x^2 - a^2)^2 - t)(x^2 - t^k)$, $a \neq 0$; (4) $((x^2 - a^2)^2 - t)(x^2 - b(t))$, $b(t) \neq 0$. If F_2 is of type (1), then D is irreducible and smooth. As a smooth triple cover of $C \cong \mathbf{P}^1$, the ramification index of D is at least 4. Hence F_3 is of type (1) as well. Let $D \sim 3C'_0 + aF'$. Then $2DC' = D(D + K_{Q/C}) = 4$, impossible. If F_2 is of type (2), then D is smooth and cannot be irreducible. D has 2 components $D_1 \sim 2C'_0 + aF'$ and $D_2 \sim 2C'_0 + bF'$. By the same argument, we have $D_1C' = D_1(D_1 + K_{Q/C}) + 2$. Hence $D_1C'_0 = 0$ and $D_1D_2 = 0$. It is impossible. $\square$

Suppose that G is a cyclic automorphism group of f . Similarly, there is an exact sequence

$$1 \longrightarrow K \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 1$$

where $H \subseteq \text{Aut}(C)$, $K = \{(\tilde{\sigma}, \text{id}) \in G\}$. It is known that $|K| \leq 10$.

Lemma 4.2 *Suppose that $f : S \longrightarrow C$ is a fibration and that G is a cyclic automorphism group of f . If there exists a point $p \in C$ such that*

- (1) $\sigma|_{f^{-1}(p)} \in K|_{f^{-1}(p)}$, for $\sigma \in G$ and σ stabilize $f^{-1}(p)$;
- (2) $K \longrightarrow \text{Aut}(f^{-1}(p))$ is injective.

Then $|K|$ and $|\text{Stab}_H(p)|$ are coprime.

Proof. Let $H_1 = \text{Stab}_H p$, $F = f^{-1}(p)$. Let $h = |H_1|$, $k = |K|$, $d = (h, k)$. Assume that σ is a generator of $\beta^{-1}(H_1)$. Then $\beta((\sigma^{k/d})^h) = 1$ implies $\sigma^{hk/d} \in K$. On the other hand, since $\sigma|_F \in K|_F$ by (1), we obtain $(\sigma^{h/d})^k|_F = \text{id}_F$. Thus $\sigma^{kh/d} = 1$ by (2). This is impossible. $\square$

Proposition 4.3 *If S is a surface of general type which has a relatively minimal fibration of genus two $f : S \longrightarrow C$ with $g(C) = 1$, G is a cyclic automorphism group of f , then*

$$|G| \leq 5K_S^2$$

for $K_S^2 \geq 12$.

Proof. If H does not act freely on C , then $|H| \leq 6$. Hence $|G| \leq 60$ and the conclusion holds. Therefore we will assume H acts freely afterwards. So $G \cong K \times H$ and G is cyclic if and only if $(|K|, |H|) = 1$. We will discuss case by case.

Case I. Suppose that the horizontal part R' of the branch locus R is étale over C . There exists a singular fiber F_0 with $|\text{Stab}_H f(F_0)| = 1$. It is not difficult to show that in this case

$$|G| \leq 5K_S^2.$$

Case II. Suppose that R' is not étale.

(a) $\bar{K} \cong Z_5$. Let F_0 be a singular fiber. The local equation of R' near F_0 is (1) $x(x^5 - t^k)$ or (2) $x(t^k x^5 - 1)$, $k = 1, 2$. We construct the quotient surface $Q = P/\bar{K}$ as in Section 3. R' must contain one of the section C_0 or C_∞ . We take this section away from R' , get a reduced divisor R_1 with $R_1 F = 5$. Let $D = \alpha(R_1)$, then $D \sim C'_0 + aF'$. Since $DC'_0 = 0$, $a = e' = 5e$. Thus $R_1 \sim 5C_0 + 5eF$ and $R_1 C_\infty = 5e$. Since the intersection number of R_1 and F on the fiber F_0 is equal to $k \leq 2$, the number of singular fibers must be a multiple of 5. But $|H|$ can not be divided by 5, hence the singular fibers are located in different H -orbits. This means $|G| \leq 5K_S^2$.

(b) $\bar{K} \cong Z_4$. The local equation of R' near a singular fiber F_0 is $x(x^4 - t^k)$, $k = 1, 2$. We use the same construction as in case (a). Then R' must contain C_0 and C_∞ . Let $R_1 = R' - C_0 - C_\infty$, $D = \alpha(R_1)$. Then $D \sim C'_0 + e'F'$. Similarly we deduce $R_1 C_\infty = 4e$. Since $|H|$ cannot be even, there are more than one singular H -orbits. So $|G| \leq 5K_S^2$.

(c) $\bar{K} \cong Z_3$. If f has only one H -orbit of singular fibers and that $|G| > 5K_S^2$, then $s_2(F_0) = 5$, namely, the local equation of R' is $x^6 + a(t)x^3 + t$. Constructing the quotient surface $Q = P/\bar{K}$, $D = \alpha(R') \sim 2C'_0 + aF'$ is a smooth irreducible curve and $r \neq |H|$. Since $DC'_0 = 0$, $DC'_\infty = |H|$, we get $a = e' = 3e = |H|$, i.e. $(|H|, |K|) = 3$, a contradiction.

(d) $\bar{K} \cong Z_2$. Lemma 3.7 ensures $|K| \leq 5K_S^2$.

(e) $\bar{K} = 1$. If $s_2(F_0) \geq 2$, then $|G| \leq 5K_{S/C}^2$. If $s_2(F_0) = 1$, there is only one situation, i.e. the local equation of R' near F_0 is $(x^2 - t)(x - a_1(t))(x - a_2(t))(x - a_3(t))(x - a_4(t))(x - a_5(t))$, $a_i(0) \neq 0$. Suppose that there is only one singular H -orbit. Then R' is a smooth 6-tuple cover of C . The contribution of each singular fiber to the ramification index equals 1. By Hurwitz formula,

$$2g(R') - 2 = 6(2g(C) - 2) + |H|.$$

So $|H|$ is even, a contradiction. $\square$

Proposition 4.4 *If S is a surface of general type which has a relatively minimal fibration of genus two $f : S \longrightarrow C$ with $g(C) = 0$, G is a cyclic automorphism group of f , then*

$$|G| \leq 12.5K_S^2 + 90.$$

Proof. If R' is étale, we have $|G| \leq 5K_{S/C}^2$. If there is a singular fiber in a big H -orbit with $s_3(F) > 0$, then $|G| \leq (50/7)K_{S/C}^2$. Now assume that R' has only negligible singularities or ramifications in big H -orbits. If $\bar{K} \cong Z_4$ or 1, we have $|G| \leq 10K_{S/C}^2$ by Lemma 2.8. When $\bar{K} \cong Z_3$ or Z_2 , if f has only one H -orbit of singular fibers, then Lemmas 3.6 and 3.7 ensure $|G| \leq 6K_{S/C}^2$. Otherwise, by the proof of Proposition 4.2, f has at least 2 big H -orbits of singular fibers, hence $|G| \leq 10K_{S/C}^2$.

It remains the case of $\bar{K} \cong Z_5$. The proof of Proposition 4.3 tells us that if f has only one big H -orbit of singular fibers, then f has another singular fiber which is stabilized by H . By Lemma 2.4, we have

$$K_{S/C}^2 \geq \frac{4}{5}(|H| + 1),$$

so

$$|G| + 10|H| \leq 12.5K_{S/C}^2 - 10 = 12.5K_S^2 + 90. \quad \square$$

When $g(C) \geq 2$, we need the following lemma on the order of some automorphisms of a curve. The proof of the lemma is just a slight modification of that of the theorem of Wiman[5]. For the convenience of the reader, we include its proof here which is a modified copy of the version given in [8, Lemma B].

Lemma 4.3 *Let H be a cyclic group of automorphisms of a curve C of genus $g \geq 2$ such that the order of $|Stab_H(p)|$ is odd for any $p \in C$. Then*

$$|H| \leq 3g + 3.$$

Proof. Let x be a non-zero element in H with maximal number of fixed points, H' the subgroup of H generated by elements fixing all fixed points of x , n the number of fixed elements of x , k the order of H' . Then k must be

odd. Let $C' = C/H'$, $g' = g(C')$, and let Σ be the image of the set of fixed points of H' on C' . We have

$$2g - 2 = 2kg' - 2k + n(k - 1). \quad (1)$$

and the quotient group $H'' = H/H'$ is a cyclic group of automorphisms of C' which satisfies the same condition imposed on H , i.e. $|\text{Stab}_{H''}(p)|$ is odd for any $p \in C'$.

If $n = 0$, then $g' \geq 2$ and $|H| \leq g - 1$. If $n = 2$, then because every non-zero element of H'' induces a non-trivial translation on Σ , we must have $|H''| \leq 2$, so $|H| \leq 2k$, then $|H| \leq 2g$ by (1) (note that $g' \neq 0$ in this case). So we may assume $n \geq 3$.

Suppose $g' = 1$, H'' acts freely on C' . Considering the induced action H'' on Σ , we see that $|H''| \leq n$. So (1) gives $|H| \leq 2g + n - 2$. On the other hand, since $k \geq 3$, (1) also gives $n \leq g - 1$, therefore

$$|H| \leq 3g - 3$$

in this case.

Suppose $g' = 1$, H'' does not act freely on C' , then H'' has a fixed point. By assumption, $|H''|$ must be odd. This implies $|H''| \leq 3$. So (1) gives

$$|H| \leq 2g + 1.$$

Now suppose that C' is a rational curve. Then the action of H'' has exactly 2 fixed points. So $|H''|$ must be odd. If one of these two points is in Σ , then $|H''| \leq n - 1$ in view of the action of H'' on Σ . Since $|H''|$ is odd, we have $n \geq 4$. So

$$|H| \leq 3g + 3.$$

Suppose that Σ and the two fixed points, ξ, η of H'' are disjoint. Let $H_1 \subset H$ be the stabilizer of a point in the inverse image of ξ . Then $[H : H_1] = k$. As the stabilizer of a point in the inverse image of η is also of index k in H , we see that any non-zero element in H_1 fixes exactly $2k$ points, i.e., the inverse image of ξ and η . Now we can replace H' by H_1 and repeat the arguments above (note that the only conditions we used are that non-trivial elements in H' have same fixed point set and that H/H' acts faithfully on Σ). But then Σ is composed of two orbits of H'' , so $|H''| \leq n/2$, whereby

$$|H| \leq \frac{3}{2}g + 3$$

by (1).

At last we use induction on g . Suppose that $g' \geq 2$ and $|H''| \leq 3g' + 3$. (1) gives

$$3g + 3 - (n - 4) \frac{3(g - g')}{2g' - 2 + n} \geq |H|.$$

If $n \geq 4$, we have done. If $n = 3$, by assumption, we must have $|H''| \leq 3$. Therefore

$$|H| \leq \frac{3(2g + 1)}{2g' + 1} \leq \frac{3}{5}(2g + 1) \leq 3g + 3. \quad \square$$

Proposition 4.5 *If $f : S \longrightarrow C$ is a relatively minimal fibration of genus 2 with $g(C) \geq 2$, G is a cyclic automorphism group of f , then*

$$|G| \leq 5K_S^2 + 30$$

for $K_S^2 \geq 48$.

Proof. (1) Assume that $|H| = 4g(C) + 2$ and $|K| = 10$. Let $g = g(C)$. By the theorem of Wiman (see the version given in [8, Lemma B]), C is a cyclic cover of $\mathbf{P}^1$ with ramification index $r_1 = 2$, $r_2 = 2g + 1$, $r_3 = 4g + 2$ or $r_1 = 3$, $r_2 = 6$, $r_3 = (4g + 2)/3$. In fact, these r_i are the orders of $\text{Stab}_H(p)$ for $p \in C$. Since Z_{10} is a maximal cyclic automorphism subgroup of a smooth curve of genus 2, by Lemma 4.2 we have $(|\text{Stab}_H(p)|, |K|) = 1$ if $f^{-1}(p)$ is a smooth fiber. But in case 1, r_1 and r_3 are even, in case 2, r_2 and r_3 are even. So f has at least $(2g + 10)/3$ singular fibers. By Lemma 2.4, we have $s_2(F) \geq 4$ for a singular fiber F . Hence

$$K_S^2 - 8(g - 1) = K_{S/C}^2 \geq \frac{4}{5} \cdot \frac{2g + 10}{3} = \frac{8(g + 5)}{15},$$

$$|G| = 10|H| = 40g + 20 \leq \frac{75}{16}K_S^2 + 45 \leq 5K_S^2 + 30$$

when $K_S^2 \geq 48$.

If $|K| \leq 8$ and $|K|$ is even, then by Lemma 4.3 there exist points $p \in C$ with $(|\text{Stab}_H(p)|, 2) \neq 1$. Hence $K_S^2 - 8(g - 1) = K_{S/C}^2 \geq 1$ and

$$|G| \leq 8|H| = 32g + 16 \leq 4K_S^2 + 44 \leq 5K_S^2 + 30$$

when $K_S^2 \geq 14$.

If $|K|$ is odd, then $|K| \leq 5$. The inequality is immediate.

(2) Assume that $|H|$ is odd. By Lemma 4.3, we have

$$|H| \leq 3g + 3.$$

So

$$|G| \leq 10|H| \leq 30g + 30 \leq \frac{15}{4}K_S^2 + 60 \leq 5K_S^2 + 30$$

when $K_S^2 \geq 24$.

(3) Assume that $|H|$ is even and $|H| < 4g + 2$. If $|K| = 10$, f must have more than one singular fibers by Lemma 2.4. So $K_S^2 - 8(g - 1) = K_{S/C}^2 \geq 2$. We get

$$|G| = 10|H| \leq 40g \leq 5K_S^2 + 30.$$

If $|K| \leq 8$, it is not difficult to obtain this inequality. $\square$

It seems that this bound is not the best. In Section 5 we will give an example to show there are infinitely many fibrations which has an automorphism with order $3.75K_S^2 + 60$.

5 Examples

Example 5.1 *Fibration with $|G| = 50K_S^2$.*

Let C be a Hurwitz curve, i.e. $|\text{Aut}(C)| = 84(g(C) - 1)$, F be a curve of genus 2 with $|\text{Aut}(F)| = 48$. Let $S = C \times F$, $f = p_1 : S \rightarrow C$. Then $K_S^2 = 8(g(C) - 1)$, $\text{Aut}(f) \cong \text{Aut}(C) \times \text{Aut}(F)$,

$$|\text{Aut}(f)| = |\text{Aut}(C)| \cdot |\text{Aut}(F)| = 504K_S^2.$$

Example 5.2 *Fibrations with $|G| = 126K_S^2$ which is not locally trivial.*

Let $F = \mathbf{P}^1$. Let $p_1 = 0$, $p_2 = \infty$, $p_3 = 1$, $p_4 = \sqrt{-1}$, $p_5 = -1$, $p_6 = -\sqrt{-1}$ be 6 points on F . Let C be a Hurwitz curve. Then C has an H -orbit $\{q_1, \dots, q_m\}$ which contains $m = 12(g(C) - 1)$ points. Let $P = C \times F$. Taking $R = p_1^*(q_1 + \dots + q_m) + p_2^*(p_1 + \dots + p_6)$ as the branch locus, we construct a double cover of P . After desingularization, we get a smooth surface S with a genus 2 fibration $f : S \rightarrow C$. By computation, we obtain $K_S^2 = 32(g(C) - 1)$, $|G| = 48 \times 84(g(C) - 1) = 126K_S^2$.

Example 5.3 *Fibrations with $|G| = 144K_S^2$ and $g(C) = 1$.*

Let F and $p_1, \dots, p_6$ as Example 5.2. Let C be an elliptic curve with j -invariant $j(C) = 0$. Fix a $q_1 \in C$, then the order of the group of automorphisms $\text{Aut}(C, q_1)$ of C leaving q_1 fixed is equal to 6. Let $H_1 \cong Z_m \oplus Z_m$ be a subgroup of translations of $\text{Aut}(C)$. Take an extension subgroup $H_1 \subset H \subset \text{Aut}(C)$ such that $H/H_1 \cong \text{Aut}(C, q_1)$. Then $|H| = 6m^2$. Let $q_1, \dots, q_{m^2}$ be the orbit of q_1 under H . Let $P = C \times F$. Using $R = p_1^*(q_1 + \dots + q_{m^2}) + p_2^*(p_1 + \dots + p_6)$ as the branch locus, we construct a double cover of P . After desingularization, we get a smooth surface S with a genus 2 fibration $f : S \rightarrow C$. By computation, we get $K_S^2 = 2m^2$. On the other hand, $|K| = 48$ gives $|G| = 288m^2 = 144K_S^2$.

Example 5.4 *Rational fibration with $|G| = 120(K_S^2 + 8)$.*

Let F and $p_1, \dots, p_6$ as Example 5.2. Let $C = \mathbf{P}^1$, $q_1, \dots, q_{12}$ be the 12 vertices of an icosahedron. Let $P = C \times F$. Taking $R = p_1^*(q_1 + \dots + q_{12}) + p_2^*(p_1 + \dots + p_6)$ as the branch locus, we can construct a double cover of P . After desingularization, we obtain a genus 2 fibration $f : S \rightarrow C$ with $K_S^2 = 16$, $|H| = 60$, $|K| = 48$, $|G| = 2880 = 120(K_S^2 + 8)$.

Example 5.5 *Rational fibrations with $|G| = 48(K_S^2 + 8)$.*

Let F and $p_1, \dots, p_6$ as Example 5.2. Let $C = \mathbf{P}^1$, $q_1, \dots, q_m$ be the m -th roots of unit. Then use the same construction as Example 5.2, we obtain a genus 2 fibration with $K_S^2 = 2(m - 4)$, $|K| = 48$, $|H| = 2m$, $|G| = 96m = 48(K_S^2 + 8)$.

Example 5.6 *Exceptional rational fibrations listed in the proof of Proposition 3.4.*

Using the same construction as Example 5.2, take $q_1, \dots, q_{20}$ as the 20 vertices of a dodecahedron. We get a fibration with $K_S^2 = 32$, $|G| = 2880 = 90K_S^2$. If we take $q_1, \dots, q_6$ as the 6 vertices of an octahedron, we get a fibration with $K_S^2 = 4$, $|G| = 1152 = 288K_S^2$. If we take $q_1, \dots, q_8$ as the 8 vertices of a cube, we get a fibration with $K_S^2 = 8$, $|G| = 1152 = 144K_S^2$.

Example 5.7 *Fibrations the order of whose abelian automorphism group is $12.5(K_S^2 + 8)$.*

Let $x_0, \dots, x_{2m}, x_{2m+1}$ be the homogeneous coordinates in $\mathbf{P}^{2m+1}$, $\mathbf{P}^{2m}$ be the hyperplane defined by $x_{2m+1} = 0$. Let $\varphi : t \mapsto (1, t, \dots, t^{2m}, 0)$ be a $2m$ -uple embedding of $\mathbf{P}^1$ in $\mathbf{P}^{2m}$ and denote its image by Y . Then Y is a rational normal curve of degree $2m$. Let X be the cone over Y in $\mathbf{P}^{2m+1}$ with vertex $P_0(0, 0, \dots, 0, 1)$. Denote $\eta = \exp(2\pi i/10m)$. Then the automorphism $\sigma : (x_0, \dots, x_{2m+1}) \mapsto (x_0, x_1\eta, \dots, x_{2m}\eta^{2m}, x_{2m+1})$ of $\mathbf{P}^{2m+1}$ is of order $10m$. The automorphism $\tau : (x_0, \dots, x_{2m+1}) \mapsto (x_0, \dots, x_{2m}, x_{2m+1}\eta^{2m})$ of $\mathbf{P}^{2m+1}$ is of order 5. The cone X is stabilized by these automorphisms σ and τ . Take a hypersurface H defined by $x_0^5 + x_{2m}^5 + x_{2m+1}^5$ which is also stabilized by σ and τ . Moreover, $P_0 \notin H$. Now blowing up the cone X at the vertex P_0 we get a Hirzebruch surface $P = F_{2m}$ which has an automorphism $\tilde{\sigma}$ of order $10m$ induced by σ and an automorphism $\tilde{\tau}$ of order 5 induced by τ . The pull-back of the intersection $H \cap X$ is a smooth divisor R_1 on P which is linearly equivalent to $5C_0 + 10mF$. Taking $R = R_1 + C_0 \equiv 6C_0 + 10mF$ which is a smooth even divisor and stabilized under $\tilde{\sigma}$ and $\tilde{\tau}$, as the branch locus, we can construct a double cover S of P which has a natural genus 2 fibration $f : S \rightarrow \mathbf{P}^1$. Since $K_P \equiv -2C_0 - (2m+2)F$, $K_S^2 = 2(K_P + R/2)^2 = 8(m-1)$. The pull-back of $\tilde{\sigma}$ on S can generate a cyclic automorphism subgroup H of order $10m$. The pull-back of $\tilde{\tau}$ on S together with the hyperelliptic involution of the fibration f generates a cyclic automorphism subgroup $K \cong Z_{10}$. As H and K commute, $G = KH \cong Z_{10} \oplus Z_{10m}$ is an abelian automorphism group of f with order $|G| = 100m = 12.5(K_S^2 + 8)$.

Example 5.8 *Rational fibrations which has an automorphism with order $12.5K_S^2 + 90$.*

Let $x_0, \dots, x_{2m}, x_{2m+1}$ be the homogeneous coordinates in $\mathbf{P}^{2m+1}$, $\mathbf{P}^{2m}$ be the hyperplane defined by $x_{2m+1} = 0$. Let $\varphi : t \mapsto (1, t, \dots, t^{2m}, 0)$ be a $2m$ -uple embedding of $\mathbf{P}^1$ in $\mathbf{P}^{2m}$ and denote its image by Y . Then Y is a rational normal curve of degree $2m$. Let X be the cone over Y in $\mathbf{P}^{2m+1}$ with vertex $P_0(0, 0, \dots, 0, 1)$. Denote $\eta = \exp(2\pi i/(50m-5))$. Then the automorphism $\sigma : (x_0, \dots, x_{2m+1}) \mapsto (x_0, x_1\eta^5, \dots, x_{2m}\eta^{10m}, x_{2m+1}\eta)$ of $\mathbf{P}^{2m+1}$ is of order $50m-5$. The cone X is stabilized by this automorphism σ . Take a hypersurface H defined by $x_0^4x_1 + x_{2m}^5 + x_{2m+1}^5$ which is also stabilized by σ and $P_0 \notin H$. Now blowing up the cone X at the vertex P_0 we get a Hirzebruch surface $P = F_{2m}$ which has an automorphism $\tilde{\sigma}$ of order $50m-5$ induced by σ . The pull-back of the intersection $H \cap X$ is a

smooth divisor R_1 on P which is linearly equivalent to $5C_0 + 10mF$. Taking $R = R_1 + C_0 \equiv 6C_0 + 10mF$ which is a smooth even divisor and stabilized under $\tilde{\sigma}$, as the branch locus, we can construct a double cover S of P which has a natural genus 2 fibration $f : S \longrightarrow \mathbf{P}^1$. Since $K_P \equiv -2C_0 - (2m+2)F$, $K_S^2 = 2(K_P + R/2)^2 = 8(m-1)$. The pull-back of $\tilde{\sigma}$ on S can generate a cyclic automorphism group G_1 of order $50m-5$. Since $|G_1|$ is odd, G_1 and the hyperelliptic involution of the fibration f generate a cyclic automorphism group G of S . Therefore $|G| = 100m - 10 = 12.5K_S^2 + 90$.

Example 5.9 *Fibrations which has an automorphism with order $5K_S^2$.*

Let $F = \mathbf{P}^1$. Let $p_1 = 0, p_k = \exp(2k\pi i/5), k = 1, \dots, 5$, be 6 points in F . Let C be an elliptic curve, $q_1, \dots, q_m$ be an orbit of a cyclic translation group $H \subseteq \text{Aut}(C)$ of order m where m is an odd prime different from 5. Then using the same construction as Example 5.2, we obtain a genus 2 fibration with $K_S^2 = 2m$, $K \cong Z_{10}$. Let $G = K \times H \cong Z_{10m}$. $|G| = 10m = 5K_S^2$.

Example 5.10 *Fibrations which has an automorphism with order $3.75K_S^2 + 60$.*

Let $p_0 = 0, p_k = \exp(2k\pi i/3), k = 1, 2, 3$ be 4 points in $C' = \mathbf{P}^1$. For any odd prime $m \neq 3, 5$, taking $D = p_0 + p_1 + (m-1)p_2 + (m-1)p_3$ as a branch locus, we can construct a m -cyclic cover $\sigma : C \longrightarrow C'$. Then $g(C) = m-1$. $H'' = \{x \mapsto x \exp(2k\pi i/3) | k = 1, 2, 3\} \cong Z_3$ is a cyclic automorphism group of C' which stabilizes the set $\{p_0, p_1, p_2, p_3\}$. On the other hand, the Galois group H' of the m -cyclic cover σ is isomorphic to Z_m . We can obtain an extension

$$1 \longrightarrow H' \longrightarrow H \longrightarrow H'' \longrightarrow 1$$

such that $Z_{3m} \cong H \subseteq \text{Aut}(C)$.

Let $q_0 = 0, q_k = \exp(2k\pi i/5), k = 1, \dots, 5$, be 6 points in $F \cong \mathbf{P}^1$. Let $P = C \times F$. Take $R = p_2^*(q_0 + q_1 + \dots + q_5)$ as branch locus, we can construct a double cover $\theta : S \longrightarrow P$ which is also a genus 2 fibration $f = p_1 \circ \theta : S \longrightarrow C$. F has a cyclic automorphism group $K_1 = \{y \mapsto y \exp(2k\pi i/5) | k = 1, \dots, 5\} \cong Z_5$ which stabilizes the set $\{q_0, \dots, q_5\}$ and can be lift to P . It is not difficult to see that we can get $K \cong Z_{10}$ by adding

the involution of the double cover. Then $G = K \times H \cong Z_{30m}$ is a cyclic automorphism group of f which satisfies

$$|G| = 30m = 30(g(C) + 1) = \frac{15}{4}K_S^2 + 60,$$

because $K_S^2 = 8(g(C) - 1)$.

References

- [1] Andreotti, A., Sopra le superficie che posseggono trasformazioni birazionali in se, *Rend. Mat. Appl.* **9**(1950), 255–279.
- [2] Beauville, A., Le nombre minimum de fibres singulières d’une courbe stable sur $\mathbf{P}^1$, *Astérisque* **86**(1981), 97–108.
- [3] Horikawa, E., On algebraic surfaces with pencils of curves of genus 2, in *Complex Analysis and Algebraic Geometry*, 79–90, Cambridge Univ. Press, 1977.
- [4] Tsuji, R., On conformal mapping of a hyperelliptic Riemann surface onto itself, *Kodai Math. Sem. Rep.* **10**(1958) 127–136.
- [5] Wiman, A., Über die hyperelliptischen Kurven und diejenigen von Geschlecht $P = 3$, welche eindeutige Transformationen in sich zulassen, *Bihang Till. Kongl. Svenska Vetenskaps-Akademiens Handlingar* **21**(1895/96), 1–23.
- [6] Xiao, G., *Surfaces Fibrées en Courbes de Genre Deux*, Lecture Notes in Math. 1137, Springer-Verlag, 1985.
- [7] Xiao, G., On abelian automorphism group of a surface of general type, *Invent. Math.* **102**(1990), 619–631.
- [8] Xiao, G., On the stable reduction of pencils of curves, *Math. Z.* **203**(1990), 379–389.
- [9] Xiao, G., π_1 of elliptic and hyperelliptic surfaces, *Intern. J. of Math.* **2**(1991), 599–615.

- [10] Xiao, G., *The Fibrations of Algebraic Surfaces*, Shanghai Scientific & Technical Publishers, 1992 (in Chinese)
- [11] Xiao, G., Bound of automorphisms of surfaces of general type, I, to appear in *Annals of Math.*
- [12] Xiao, G., Bound of automorphisms of surfaces of general type, II, preprint

Department of Mathematics
East China Normal University
Shanghai 200062
People's Republic of China